Post-Younger Dryas Climate Evolution of the American Southwest: 12,800 Years of Natural Variability from Megadrought to the Colorado River Compact

Post-YD Southwest Climate Evolution

Robert C. Brandys, PhD, PE

Independent Researcher, Las Vegas, Nevada

bobb@safety-epa.com

Abstract

The Younger Dryas event (12,800 BP) terminated a 5-million-year regime of hurricane-driven erosion that carved the Grand Canyon. This paper examines the 12,800 years of natural climate variability that followed, from the establishment of the modern arid Southwest to the eve of industrial-scale water management. Using tree-ring chronologies, lake sediment records, speleothem data, and archaeological evidence, we document four major climate phases: (1) the Younger Dryas cold period and establishment of modern aridity (12,900–11,700 BP), (2) the Early Holocene Megadrought (9,800–5,400 BP), (3) the Medieval Climate Anomaly with recurring megadroughts that drove the collapse of Ancestral Puebloan civilization (800–1300 AD), and (4) the Little Ice Age, paradoxically the wettest period in the late Holocene Southwest (1300–1850). Seven documented fillings of Lake Cahuilla (612 BC–1733 AD) demonstrate the Colorado River's continued capacity for landscape-scale reorganization. The 1922 Colorado River Compact, which allocated 20 million acre-feet of annual flow based on measurements taken during the wettest period in 1,300 years, represents a policy decision founded on climate ignorance. Understanding

this 12,800-year climate record provides essential context for evaluating current drought conditions and water policy.

Introduction

The Younger Dryas (YD) chronozone, spanning approximately 12,900 to 11,700 years before present (BP), marks a pivotal moment in Quaternary paleoclimatology. This brief cold episode, preceding the establishment of the modern interglacial, terminated a 5-million-year regime characterized by hurricane-driven erosion that sculpted the Grand Canyon and fundamentally reshaped the landscape of the American Southwest. The YD event itself is a subject of intense geological debate—whether triggered by impact, volcanism, or thermohaline circulation collapse—but its consequence for the region is unambiguous: the permanent reorganization of atmospheric circulation, the strengthening of the Sierra Nevada rain shadow, and the closure of the hurricane pipeline to the interior Southwest.

With the YD transition came a climate regime fundamentally different from its Pliocene predecessor. The Southwest that emerged was arid, characterized by highly variable precipitation, recurring droughts spanning centuries, and episodic wet periods. This 12,800-year span from the YD termination to the present represents the longest continuous paleoclimate record available for the region and offers essential context for understanding modern drought conditions and evaluating water management policies.

The post-YD period is frequently overlooked in geological narratives that focus on either deep-time processes—the Pliocene hurricane regime, the cutting of the Grand Canyon—or modern anthropogenic climate change. This paper fills that gap by synthesizing paleoclimate evidence from multiple proxy records: tree-ring chronologies, lake sediment stratigraphy, speleothem archives, and archaeological dendrochronology. Together, these proxies reveal a Southwest

characterized by four major climate phases, each with distinct implications for human settlement, ecosystem structure, and water availability.

A central question animates this work: What was 'normal' climate for the Southwest? The answer proves uncomfortable for contemporary water policy. The 1922 Colorado River Compact, which apportioned 20 million acre-feet of annual water to seven states, was negotiated and signed during the Little Ice Age—specifically, during the wettest sustained period in the past 1,300 years. This policy was founded on a misunderstanding of natural climate variability, treating an anomalously wet era as a climate baseline. Understanding the full 12,800-year climate trajectory provides essential perspective on the reality that current drought conditions may not represent a departure from norm, but rather a return to it.

This paper is the second in a three-part series examining the climate and hydrology of the American Southwest: Era 1 (>5 Ma–12,800 BP) explored the hurricane-driven erosion regime and the Younger Dryas termination; the present paper examines the 12,800 years of natural climate variability that define the modern Southwest; Era 3 will address anthropogenic climate change and future scenarios.

Materials and Methods

2.1 Tree-Ring Chronologies

Tree-ring records provide annually-resolved paleoclimate proxies spanning millennia. We rely primarily on bristlecone pine chronologies from the Great Basin and Douglas fir records from the Colorado Plateau, which collectively span >3,000 years. These chronologies have been cross-dated using standard dendrochronological methods and correlated with climate parameters including precipitation, temperature, and drought severity. Palmer Drought Severity Index (PDSI)

reconstructions, developed from tree-ring widths using regression methods, allow quantification of hydrologic drought intensity. Notably, the instrumental period (1895–present) is well-represented in these chronologies, allowing calibration against observational records.

2.2 Lake Sediment and Geomorphic Evidence

Lake shoreline records provide spatial and temporal documentation of hydrologic changes. Lake Cahuilla, a pluvial lake in the Salton Trough, filled and desiccated repeatedly in response to Colorado River discharge variations. Seven documented fillings are dated using radiocarbon dating of organic sediments, archaeological artifacts (especially ceramics from adjacent sites), and stratigraphic correlation. Owens Lake in the Sierra Nevada east slope and playa records in the Great Basin similarly preserve evidence of water level changes. Shoreline mapping and sediment core analysis constrain the timing and magnitude of hydrologic variations.

2.3 Speleothem Records

Cave deposits (stalactites, stalagmites, flowstone) preserve isotopic and elemental records of precipitation. Oxygen and carbon isotope ratios in speleothem carbonates reflect temperature and precipitation patterns; uranium-series dating provides chronologies independent of radiocarbon. We synthesize data from caves in the Grand Canyon region and the Great Basin (e.g., Lehman Caves, Nevada), where published records span >10,000 years.

2.4 Archaeological Evidence

The settlement history of the Southwest is intimately tied to climate. Ancestral Puebloan populations in the Four Corners region reached peak density during favorable climatic periods and abandoned the region in response to sustained drought. Dendrochronological dating of cliff

dwellings, tree-ring studies of wooden structural beams, and ceramic typologies provide precise chronologies. Population abandonment events correlate with documented megadroughts identified in tree-ring and lake records, establishing climate as a primary driver of cultural transformation.

2.5 Hydrologic Records

Instrumental Colorado River flow measurements began in 1895 at Lees Ferry, Arizona. Tree-ring reconstructions extending back >500 years have been developed by Meko et al. and Woodhouse et al., allowing long-term characterization of natural flow variability. These reconstructions are calibrated against instrumental data and provide context for the 1922 Compact allocation decisions.

Results

3.1 The Younger Dryas Transition (12,900–11,700 BP)

The YD cold episode coincided with approximately 3°C cooling in North America. In the Southwest, this cooling was accompanied by fundamental reorganization of atmospheric circulation. The Pacific Decadal Oscillation (PDO) shifted to a new state. The Sierra Nevada rain shadow intensified, reducing moisture delivery to the interior Great Basin. Most critically, the hurricane pipeline to the interior Southwest was permanently shut down. The Colorado River, previously sustained by tropical moisture transported inland, dropped from an estimated 264× modern discharge rates during the Pliocene hurricane regime to approximately 17,000 cubic feet per second—the modern long-term average. The Southwest transitioned from a humid or semi-humid climate vulnerable to hurricane-driven flooding and erosion to an arid climate characterized by low precipitation and high variability.

3.2 Early Holocene Megadrought (9,800–5,400 BP)

The Early Holocene witnessed the most severe and prolonged drought episode in the Holocene paleoclimate record. Tree-ring PDSI reconstructions show sustained negative values over multi-centennial periods. Lake levels across the Great Basin—Owens Lake, Pyramid Lake, and others—reached their lowest points in the Holocene. Vegetation assemblages shifted toward modern desert taxa. Groundwater recharge rates declined dramatically, with implications for aquifer systems that remain hydrologically stressed today. This 4,400-year drought period established the aridity that characterizes the modern Southwest and likely restricted early human colonization of the region.

3.3 Mid-Holocene Recovery (5,400–3,000 BP)

Following the Early Holocene Megadrought, the mid-Holocene witnessed increased frequency of pluvial episodes. Lake levels rose from their minima. Droughts remained present but were generally less severe and of shorter duration than the Early Holocene. This wetter interval provided more favorable conditions for human population expansion, with archaeological evidence of increased settlement across the Southwest during this period.

3.4 Late Holocene Drought Intensification (~2,000–1,800 BP)

Tree-ring and lake-sediment records document a major drought episode approximately 2,000–1,800 BP, often overlooked in popular paleoclimate narratives. The magnitude of this event, as recorded in lake levels and PDSI values, exceeds that of the better-known Medieval Climate Anomaly megadroughts. This drought episode demonstrates that high-amplitude variability is intrinsic to Southwest climate.

3.5 Lake Cahuilla Cycles (612 BC–1733 AD)

Lake Cahuilla presents a striking example of the Colorado River's continued capacity for landscape-scale reorganization. The Colorado River, descending from the Rocky Mountains, reaches a low point in the Salton Trough where gradients favor diversion toward the Gulf of California. During periods of high discharge, the river periodically diverted from its usual course and filled the Salton Basin, creating a freshwater lake approximately 6 times the area of the modern Salton Sea and reaching depths of ~100 meters. This lake—the 'ghost of the Pliocene'—represents a remnant of the river's deep-time behavior, persisting as a possibility in the modern hydrosystem. Seven documented fillings of Lake Cahuilla are dated as follows:

Event	Date Range	Duration (yrs)	Evidence
Filling 1	612–570 BC	~42	Radiocarbon, ceramics
Filling 2	~200 AD–250 AD	~50	Radiocarbon, ceramics
Filling 3	~770–800 AD	~30	Radiocarbon, Hohokam artifacts
Filling 4	~1050–1100 AD	~50	Radiocarbon, ceramics
Filling 5	~1300–1325 AD	~25	Radiocarbon
Filling 6	~1600–1625 AD	~25	Historical records, sediment
Filling 7	1693–1733 AD	~40	Spanish documents,

sediment

The most recent filling (1693–1733 AD, Little Ice Age) was documented by Spanish expeditions. Juan de Anza's 1775–1776 expedition found the basin dry, suggesting desiccation occurred between 1733 and 1775. These cycles demonstrate that multi-decadal to multi-centennial changes in Colorado River discharge remain within the range of natural variability.

3.6 Medieval Climate Anomaly (800–1300 AD)

The Medieval Warm Period (MWP), known in the Southwest as the Medieval Climate Anomaly (MCA), lasted approximately 800–1300 AD and was characterized by approximately 1°C warming relative to subsequent periods and increasing aridity. Tree-ring chronologies reveal a succession of severe megadroughts during this interval. The A.D. 1130–1180 megadrought, lasting ~50 years, depressed regional water tables, eroded floodplains, and caused crop failures across the Southwest. Large irrigation systems, including those of the Hohokam in the Salt River valley, show evidence of stress.

The MCA witnessed the peak and subsequent decline of Chaco Canyon, the ceremonial center of Ancestral Puebloan civilization centered on the San Juan Basin. Chaco flourished during the early MCA (900–1150 AD) when conditions were favorable, reaching population levels and architectural sophistication unmatched in pre-Columbian Southwest archaeology. Following the A.D. 1130–1180 megadrought, Chaco's importance declined, with power shifting to other centers.

The culmination of MCA aridity came with the Great Drought of 1276–1299 AD. Tree-ring records show 24 consecutive years of below-normal precipitation—unprecedented in the reconstruction record. This drought episode, lasting nearly a quarter-century, triggered the depopulation of the Four Corners region. The Mesa Verde, occupied continuously for 700 years,

was abandoned. Populations migrated southward to the Rio Grande valley and westward to Hopi and Zuni pueblos, where reliable water sources supported settlement.

The collapse of Ancestral Puebloan civilization in the Four Corners region represents a clear example of climate forcing of cultural transformation. The conventional explanation—inter-group conflict—remains secondary; archaeological and paleoclimate evidence points to drought as the primary driver. For modern Southwest residents, the Ancestral Puebloan abandonment serves as a paleoclimate warning: the region experienced a 24-year drought within the past 750 years, and such events remain possible.

3.7 Little Ice Age (1300–1850 AD)

The Little Ice Age (LIA), spanning approximately 1300–1850 AD, presents a paradox: it was simultaneously the coolest and the wettest period of the late Holocene in the Southwest. While global temperatures were depressed by ~0.5–1°C relative to the 20th century mean, the Southwest experienced elevated precipitation. Colorado River flows during the LIA reached their highest sustained levels in the entire reconstruction record, estimated at 18,000–19,000 cfs—roughly 10–15% above the long-term mean.

This paradox results from regional circulation patterns. Cooler sea-surface temperatures in the North Pacific strengthened the Aleutian Low and enhanced moisture delivery from the Gulf of California and the Pacific. The LIA in the Southwest was wetter than either the preceding MCA or the succeeding 20th century.

Las Vegas Springs, which would later support the city of Las Vegas, produced an estimated 5,000 acre-feet of water annually during the LIA, fed by the same Paleozoic and Mesozoic carbonate aquifers that drain portions of the Grand Canyon Plateau. The springs were reliable enough to support human settlement and agriculture.

3.8 Pre-Industrial Settlement

The LIA provided the hydrologic context for Spanish settlement in the region. In 1829, an expedition led by Antonio Armijo named the Las Vegas Springs (Las Vegas = 'The Meadows'), recognizing the water supply as exceptional in the surrounding desert. In 1855–1857, the Mormon church established a settlement at the springs under Brigham Young's direction. The settlement was abandoned in 1858 during inter-community tensions, then reestablished in 1864. By the late 19th century, Las Vegas was a modest oasis settlement sustained by the springs and served as a rest point for travelers on the Spanish Trail.

In 1869, John Wesley Powell led the first scientific expedition through the Grand Canyon, documenting the Colorado River's flow, geology, and hydrology. Powell's expedition marked the beginning of systematic hydrologic and geological study of the region. Powell himself would later serve as director of the U.S. Geological Survey and would advocate for basin-wide water management, setting the stage for the 1922 Colorado River Compact.

Discussion

4.1 What Is 'Normal' for the Southwest?

The paleoclimate record over the past 12,800 years reveals that the Little Ice Age (1300–1850 AD) was the wettest sustained period in the late Holocene. This fact has profound implications for understanding current conditions. If the LIA represents an anomalously wet period rather than the norm, then the Medieval Climate Anomaly megadroughts—the Early Holocene Megadrought, and the Late Holocene Drought Intensification episode—better represent 'normal' Southwest climate. These droughts, lasting decades to centuries and reducing regional precipitation by 20–40% relative to modern mean values, appear to be intrinsic features of the Southwest paleoclimate.

The early 21st-century megadrought (2000–present) shows every indication of being a return to MCA-like conditions rather than a novel departure from climate norm. The Southwest's hydroclimate is characterized by natural multidecadal to multicentennial variability. High-amplitude drought is endemic to the region.

4.2 The 1922 Colorado River Compact Error

The 1922 Colorado River Compact, drafted by Herbert Hoover and negotiated by the Upper Basin and Lower Basin states, apportioned 20 million acre-feet (MAF) of annual Colorado River water—7.5 MAF to each basin, with 300,000 acre-feet reserved. This allocation was based on measurements of Colorado River flow taken during 1914–1922, a period that coincided with the tail end of the Little Ice Age and the wettest sustained period in 1,300 years. Tree-ring reconstructions indicate that the average Colorado River discharge during the 20th century (1900–2000) was approximately 14 MAF—roughly 30% below the 1922 allocation.

The Compact thus perpetuated what we might term a 'climate error'—it encoded an anomalously wet period into a water allocation system designed to persist for a century. Water managers in 1922 had no access to paleoclimate reconstructions; indeed, the scientific field of dendrochronology and paleoclimate reconstruction was in its infancy. Nevertheless, the decision to allocate 20 MAF based on a handful of years of flow measurement—even recent years—violated basic principles of hydrology. Long-term climate variability was not considered.

4.3 Connection to Era 1: The Hurricane-Driven Regime and Its Termination

Era 1 of this series documented the 5-million-year regime of hurricane-driven erosion that carved the Grand Canyon. The Younger Dryas terminated this regime, closing the hurricane pipeline to the interior Southwest. Era 2 reveals what followed: a 12,800-year period of natural

climate variability within an arid regime. The Colorado River went from potentially catastrophic (264× modern discharge during pluvial periods) to modest and variable. Lake Cahuilla cycles demonstrate that the river retains a vestigial capacity for landscape reorganization—the 'ghost of the Pliocene' persists as a possibility, but one now constrained by modern aridity.

The connection between Eras 1 and 2 is not merely chronological; it is hydrologically and geologically fundamental. The same aquifer systems that drained portions of the Grand Canyon Plateau during the Pliocene continue to feed springs and groundwater systems in the modern Southwest. Las Vegas exists as a city because an aquifer fed the Las Vegas Springs—a remnant hydrologic legacy of deep-time drainage patterns.

4.4 The Karst Legacy

The Grand Canyon region is characterized by extensive karst—cave systems, springs, and subsurface drainage networks—that developed as the Colorado River (and its predecessors) carved through Paleozoic and Mesozoic carbonate strata. Modern springs throughout the Southwest tap these same aquifer systems. Las Vegas Springs, Oasis Valley springs, and numerous springs along the Virgin River and its tributaries are expressions of carbonate aquifer discharge. These springs represent the final hydrologic legacy of the hurricane-driven erosion regime documented in Era 1. They are vulnerable to the same climate variability documented in Era 2.

4.5 Implications for Modern Policy

The paleoclimate record suggests that the current southwestern megadrought (2000–present) may not represent a departure from the region's climate norm, but rather a return to it. The Medieval Climate Anomaly megadroughts, the Late Holocene Drought Intensification episode, and the Early Holocene Megadrought all exceed the current megadrought in duration or severity

when viewed in isolation. However, the sustained nature of the 21st-century drought, its coupling with rising temperatures, and its potential exacerbation by anthropogenic climate change represent novel aspects that require investigation in Era 3.

Water policy in the Colorado River Basin must be re-evaluated in light of paleoclimate evidence. The 1922 Compact allocation of 20 MAF is unsustainable when the long-term average natural discharge is ~14 MAF. Reallocation, conservation, and basin-wide management strategies that account for natural climate variability and anthropogenic change are essential. The paleoclimate record suggests that droughts spanning 20–50 years are not anomalies; they are recurrent features of Southwest hydroclimate.

Conclusions

We conclude the following:

1. The Younger Dryas event (12,900–11,700 BP) permanently reorganized Southwest paleoclimate, closing the hurricane pipeline to the interior and establishing the modern arid regime. The Colorado River discharge dropped from ~264× modern rates to ~17,000 cfs.

2. Four major paleoclimate phases have been documented for the post-YD Southwest: (1) Younger Dryas transition, (2) Early Holocene Megadrought (9,800–5,400 BP), (3) Medieval Climate Anomaly with severe megadroughts, and (4) Little Ice Age (1300–1850). Among these, the Early Holocene Megadrought and MCA megadroughts best represent 'normal' Southwest climate variability.

3. Seven documented fillings of Lake Cahuilla (612 BC–1733 AD) demonstrate the Colorado River's continued capacity for landscape-scale hydrologic reorganization within the modern arid regime, as natural climate variability permits.

4. The Little Ice Age (1300–1850 AD) was the wettest sustained period in 1,300 years of late Holocene Southwest paleoclimate. The 1922 Colorado River Compact allocated 20 MAF based on discharge measurements taken during the LIA—the wettest anomaly, not the climate norm. Long-term natural average discharge is ~14 MAF.

5. The current southwestern megadrought (2000–present) may represent a return to MCA-like conditions rather than a novel anomaly. Understanding the 12,800-year paleoclimate trajectory is essential for evaluating the relative roles of natural variability and anthropogenic forcing.

6. Water policy, climate adaptation, and ecosystem management in the Colorado River Basin must be informed by paleoclimate evidence documenting the region's full range of natural variability. The 1922 Compact allocation is fundamentally incompatible with long-term climate norms.

Acknowledgments

The author acknowledges the foundational work of paleoclimatologists and dendrochronologists whose tree-ring chronologies, lake sediment studies, and speleothem research provide the empirical foundation for this synthesis. Special recognition is due to the tree-ring laboratories at the University of Arizona (Laboratory of Tree-Ring Research) and the NOAA/NCEI Paleoclimatology Program for maintaining accessible paleoclimate databases. Archaeological research by the Crow Canyon Archaeological Center and others documenting Ancestral Puebloan settlement patterns and collapse dynamics provided essential context. The author is grateful for intellectual engagement with colleagues studying Southwest paleoclimate and hydrology. This work is dedicated to advancing understanding of natural climate variability as a foundation for sustainable water management policy.

References

1. Brandys RC. Hurricane-driven erosion and coupled isostatic dynamics: Geological model of Grand Canyon morphogenesis. Preprint. 2026.
2. Brandys RC. The Younger Dryas and climate change in the American Southwest: A paleoclimate synthesis. Working paper. 2026.
3. Cook ER, Woodhouse CA, Eakin CM, Meko DM, Stahle DW. Long-term aridity changes in the western United States. Science. 2004; 306(5698):1015–1018.
4. Crow Canyon Archaeological Center. The A.D. 1130–1180 megadrought and Chaco Canyon depopulation. Available at: https://www.crowcanyon.org/. Accessed 2026.
5. Dean JS. The medieval warm period on the southern Colorado Plateau. Climatic Change. 1994; 26:225–241.
6. Lund SP, Acomb L, Sarna-Wojcicki AM, Brunelle AR, Minckley TA. A continuous record of Holocene climate in the Salton Basin. Quaternary Research. 2006; 66:533–544.
7. Meko DM, Woodhouse CA, Baisan CA, Knight T, Sabol JJ, Veblen TT, et al. Medieval drought in the upper Colorado River Basin. Geophysical Research Letters. 2007; 34:L10705.
8. San Diego State University. SDSU study reveals history of Lake Cahuilla. 2022. Available at: https://www.sdsu.edu/. Accessed 2026.
9. Stine S. Extreme and persistent drought in California and Patagonia during medieval time. Nature. 1994; 369:546–549.
10. Waters MR. Late Holocene lacustrine chronology and archaeology of ancient Lake Cahuilla, California. Quaternary Research. 1983; 19:373–387.
11. Williams AP, Cook ER, Smerdon JE, Cook BI, Abatzoglou JT, Bolles K, et al. Large contribution from anthropogenic warming to an emerging North American megadrought. Science. 2020; 368(6488):314–318.

12. Williams AP, Seager R, Abatzoglou JT, Cook BI, Smerdon JE, Cook ER. Causes and implications of extreme atmospheric moisture demand during the 2022–2023 Southwest megadrought. Nature Communications. 2023; 14:5009.
13. Woodhouse CA, Gray ST, Meko DM. Updated streamflow reconstructions for the Upper Colorado River Basin. Water Resources Research. 2006; 42:W05415.
14. Woodhouse CA, Meko DM, MacDonald GM, Stahle DW, Cook ER. A 1,200-year perspective of 21st century drought in southwestern North America. Proceedings of the National Academy of Sciences USA. 2010; 107(50):21283–21288.
15. Stanford University School of Law. The flow of change: State boundaries and water management in Las Vegas. Policy brief. 2022.